

# Quotation Request 2021/22

FLUCELVAX  
TETRA

INFLUENZA VACCINE (SURFACE ANTIGEN,  
INACTIVATED, PREPARED IN CELL CULTURES)

If you wish to receive a FLUCELVAX® TETRA quotation, please complete this form and email it back to [flu.salesuk@seqirus.com](mailto:flu.salesuk@seqirus.com) or alternatively call **03450 093804**. Please be advised that this is not an order form, but a request for a quotation of your proposed order. A member of our team will contact you shortly after receiving your completed form.

Vaccine name	Licensed for	NHS List Price per dose (excl. VAT)	Number of doses required
FLUCELVAX® TETRA INFLUENZA VACCINE (SURFACE ANTIGEN, INACTIVATED, PREPARED IN CELL CULTURES)	Prophylaxis of influenza in adults and children from 2 years of age	£9.94	<input type="text"/>

## CONTACT INFORMATION (PLEASE USE CAPITAL LETTERS WHEN COMPLETING THIS FORM):


<b>Account Name</b> <input type="text"/>	<b>Contact Name</b> <input type="text"/>
<b>Address</b> <input type="text"/>	<b>Best day/time to contact</b> <input type="text"/>
<b>Town</b> <input type="text"/>	<b>Other information</b> <input type="text"/>
<b>Postcode</b> <input type="text"/>	<b>NHS practice code</b> <input type="text"/>
<b>Telephone</b> <input type="text"/>	<b>Date (DD-MMM-YYYY)</b> <input type="text"/>
<b>Email Address</b> <input type="text"/>	

By completing the quotation form, you hereby give Seqirus (and their representatives) permission to contact you.

Copyright © Seqirus UK Limited 2020.  
Date of preparation: October 2020 UK/XQIV/0720/0055(1)



## PRESCRIBING INFORMATION

**Flucelvax® Tetra**  **Influenza vaccine (surface antigen, inactivated, prepared in cell cultures)**, suspension for injection in pre-filled syringe. **Presentation:** Each 0.5ml of Flucelvax Tetra contains 15 micrograms of each of four purified virus strains propagated in Madin Darby Canine Kidney (MDCK) cells that comply with the World Health Organization quadrivalent vaccine recommendations (Northern Hemisphere) for the current season. **Indications:** Prophylaxis of influenza in adults and children from 2 years of age. **Dosage and Administration:** Adults and children aged 2 years and over should receive a single 0.5ml dose, children less than 9 years of age who have not been previously vaccinated against influenza, should receive a second dose at least 4 weeks apart. For intramuscular injection only. The preferred site for injection is the deltoid muscle of the upper arm. Young children with insufficient deltoid mass should be vaccinated in the anterolateral aspect of the thigh. **Contraindications:** Hypersensitivity to the active substance, to any of the excipients (sodium chloride, potassium chloride, magnesium chloride hexahydrate, disodium phosphate dihydrate, potassium dihydrogen phosphate), or to possible trace residues (beta-propiolactone, cetyltrimethylammonium bromide, and polysorbate 80). **Warnings and Precautions:** In order to improve the traceability of biological medicinal products, the name and the batch number of the administered product should be clearly recorded. Appropriate medical treatment and supervision should be readily available in case of an anaphylactic event following administration. Do not inject intravenously, subcutaneously or intradermally. Flucelvax Tetra must not be mixed with other vaccines in the same syringe. Vaccination should be postponed in patients with acute febrile illness until fever is resolved. As with all injectable vaccines, Flucelvax Tetra must be administered with caution to individuals with thrombocytopenia or a bleeding disorder since bleeding may occur following intramuscular administration. Syncope (fainting) can occur following or before any vaccination as a psychogenic response to the needle injection. This can be accompanied by several neurological signs such as transient visual disturbance, paraesthesia, and tonic-clonic limb movements during recovery. It is important that procedures are in place to avoid injury from faints. Endogenous or iatrogenic immunosuppression may result in insufficient antibody response. A protective immune response may not be elicited in all vaccine recipients. **Interactions:** No clinical data on co-administration with other vaccines are available. Based on clinical experience with cell-based trivalent influenza vaccine (TIVc), Flucelvax

Tetra can be given at the same time as other vaccines. **Pregnancy and Lactation:** There are limited data from the use of Flucelvax Tetra in pregnant women. However, inactivated influenza vaccines can be used at all stages of pregnancy. It is unknown whether Flucelvax Tetra is excreted in human milk. No effects on breast-fed newborn/infant are anticipated. Flucelvax Tetra may be used during lactation. **Effects on Ability to Drive and Use Machines:** Flucelvax Tetra has no or negligible influence on the ability to drive and use machines. **Side Effects:** The most common reactions ( $\geq 1/10$ ) are injection site pain, ecchymosis, erythema, induration, headache, fatigue, myalgia, loss of appetite. Additionally, injection site tenderness, irritability, sleepiness and change in eating habits were reported in children 2 to <6 years. Commonly reported adverse reactions ( $\geq 1/100$  to <1/10) include nausea, vomiting, diarrhoea, arthralgia, chills/shivering and fever ( $\geq 38^{\circ}\text{C}$ ). Vomiting in the elderly, and fever in adults and elderly were uncommon. The following have been reported post-marketing: extensive swelling of injected limb, allergic reactions (including anaphylactic shock), paraesthesia, and generalised skin reactions (including pruritus, urticaria, or non-specific rash). Paediatric subjects generally reported higher rates of local and systemic reactions compared to adults aged 18 years and over. **Overdose:** There are no data for overdose with Flucelvax Tetra.

**Legal Category:** POM. **Package Quantities:** Packs of 1 or 10 pre-filled syringes. **Marketing Authorisation Numbers:** EU/1/18/1326/001, EU/1/18/1326/002, EU/1/18/1326/003, EU/1/18/1326/004. **Basic NHS Cost:** £9.94 per 0.5ml pre-filled syringe, £99.40 per 10-pack. **Marketing Authorisation Holder:** Seqirus Netherlands B.V., Paasheuvelweg 28, 1105BJ, Amsterdam, Netherlands.

For full prescribing information and details of other side effects see the Summary of Product Characteristics at [www.medicines.org.uk/emc](http://www.medicines.org.uk/emc)

Adverse events should be reported. Reporting forms and information can be found at [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard). Adverse events relating to Seqirus products should also be reported to Seqirus UK Limited on **01748 828816**.

UK/XQIV/0720/0057(2) Date of preparation: October 2020.

Flucelvax® Tetra is a registered trademark of Seqirus UK Limited.